

HiroCyst

D-chiro-inositol 500 mg

Capsules

Composition :

Each Capsules Contains:
D-chiro-inositol 500 mg.
Standardized extract

Mfg. Lic. No. : HBL-018
D.A.R. No. : 018-31-094



**May The Best For PCOS
Treatment In Restoring
Ovulation & Normalizing
Hormone Levels**

Manufactured By :



IBN HAISAM
Herbal & Nutraceuticals
Uttarkhan, Dhaka, Bangladesh.

HyoCyst

D-chiro-inositol 500 mg

Description :

D-chiro-inositol is extracted by a patented process from Carob (*Ceratonia siliqua*) pod that mostly found in Southern Europe. Deficiency of D-chiro-inositol in peripheral tissue causes insulin resistance that leads to hyperandrogenism in ovaries which results polycystic ovary syndrome (PCOS). D-chiro-inositol helps to regulate the balance between androgen and estrogen by decreasing luteinizing hormone and serum testosterone level. It improves glucose tolerance and reduces circulating insulin. It reduces oxidative stress on follicle and decreases total cholesterol and triglyceride levels. D-chiro-inositol has been clinically proven as an effective treatment of ovarian cyst and PCOS associated metabolic disorders. It helps to improve and regulate normal ovulation after two weeks of administration. It also reduces hyperandrogenism related skin disorders like hirsutism, alopecia, acanthosis nigricans and acne. It lowers cardiovascular risk in women who are suffering from PCOS by controlling arterial pressure.


Indications and Uses:

HyoCyst capsule is indicated for the treatment and prevention of-

- * Polycystic ovary syndrome (PCOS)
- * Anovulation
- * Menstrual disorders such as amenorrhea
- * Hyperandrogenism complaints such as hirsutism, alopecia, acanthosis nigricans, acne
- * Ovarian hyperthecosis


A Promising Inositol in PCOS & Gestational Diabetes

Ovulation rate in PCOS Patient



Increased ovulation rate helps to normalize menstrual cycle and increases pregnancy rate

Plasma insulin (fasting state)



D-Chiro-Inositol decreases plasma insulin level in PCOS patient and minimized the risk of Type-2 Diabetes and Gestational Diabetes.

Dosages & Administration : 1-2 Capsules daily or as directed by the physician.

Duration of Use: Duration of the treatment should be minimum 2 months.

Contraindication: Contraindicated in patient with known hypersensitivity to D-chiro-inositol.

Pregnancy and Lactation: Several human studies have been conducted with inositol has no clinically relevant adverse effects.

Side Effects: Well tolerated in recommended dose.

Drug Interactions: Concurrent use with digoxin may result in increased digoxin toxicity.

Storage: Store at temperature of below 30 C, protect from light & moisture.

Storage : Keep in cool, dry place & keep away from the direct sunlight. Keep the medicine out of the reach of children.

How Supplied : Each box contains 15 x 1 =15 Capsules in blister pack.